



Prior To Surgery

- Continue Prescription Blood Thinners or Aspirin therapy.
- The Surgical Center will call you to let you know the exact time to arrive for your procedure. They will also advise you on which of your current medications you should take prior to surgery.
- DO NOT EAT OR DRINK anything after midnight prior to surgery unless instructed by the Surgical Center.
- Review your drop schedule carefully prior to your surgery. You must start Ofloxacin 3 days prior to surgery.
- When applying drops, allow 5 minutes between each drop to insure they are properly absorbed.

Tip: To apply drops recline back, close eye and apply drop in corner pocket of eye, then open your eye. The medication will spread across the surface of your eye.

- Make sure you start your drops in the correct eye, review your surgical schedule carefully. We will be unable to perform surgery if the drops have been placed in the wrong eye. Also, the lens type that you selected for your surgery is the lens that will be ordered for your procedure. If there is a discrepancy on the day of surgery, your surgery may be cancelled.

THE DAY OF SURGERY

- You will need a driver the day of surgery. They will need to remain at the Surgical Center during your procedure.
- Take sunglasses with you the day of surgery to wear home.
- You Do NOT need to use your drops the morning of surgery.
- Start Drops when you arrive home from Surgery (Ofloxacin & Prednisolone). Make sure you use each drop 3 times.
Example:
AM Surgery-2pm/6pm/Bedtime PM
Surgery-Arrive Home/7pm & Bedtime

- You may return to your regular diet and medications following surgery.
- The Surgical Center will provide you with a plastic shield to wear after surgery. Please wear the shield the day of surgery (except when applying your drops) and sleep in it that night. The next morning you may remove the plastic shield and only wear when you are sleeping for 5 nights.

1-Day Post-Operative Visit

- You will have a follow up appointment at the Clinton Township location the day following surgery.
- Your eye was dilated during surgery, it could take 24-48 hours for the dilation to resolve.

Contact Information

Please contact Fite Eye Center if you have any questions or concerns at 586-226-2020.

Visit our website at www.fiteeyecenter.com